Eur. Phys. J. C manuscript No.
(will be inserted by the editor)

Triple Higgs coupling in the most general 2HDM at SM-like scenario

Ginzburg I.F.[1,2]

[1]Sobolev Institute of Mathematics, av. Koptyug, 4, Novosibirsk, 630090, Russia,
[2]Novosibirsk State University, Pirogova str., 2, Novosibirsk, 630090, Russia

arXiv:1510.08270v2 [hep-ph] 23 Jul 2016

the date of receipt and acceptance should be inserted later

Abstract We consider the triple Higgs coupling for $h(125)$ Higgs boson within the most general 2HDM. At moderate values of parameters, noticeable deviation of this coupling from its SM value is improbable. This deviation can be sizable only if some measurable parameters of the model are exotic.

Keywords Higgs models · triple Higgs vertex

PACS 12.60 Fr

1 Introduction

The recent discovery of a Higgs boson with $M \approx 125$ GeV at the LHC [1]-[3] suggests that the spontaneous electroweak symmetry breaking is most probably brought up by the Higgs mechanism. The simplest realization of the Higgs mechanism introduces a single scalar isodoublet ϕ with the Higgs potential $V_H = -m^2(\phi^\dagger\phi)/2 + \lambda(\phi^\dagger\phi)^2/2$. This model is usually called "the Standard Model" (SM). The mentioned data do not rule out the possibility of realization of *beyond SM models* (BSM) which include both neutral Higgs scalars h_a (generally without definite CP parity) and charged Higgs scalars $H^{\pm b}$ with masses M_a and $M^b_\pm$ respectively.

In the discussion that follows we use the relative couplings for each neutral Higgs boson h_a (for the case with single charged Higgs boson $H^\pm$):

$$\chi^P_a = \frac{g^P_a}{g^P_{\rm SM}} \quad [P = V\,(W,Z),\ q = t,b,...,\ \ell = \tau,...]\,,$$
$$\chi^\pm_a = \frac{g(H^+H^-h_a)}{2M^2_\pm/v}, \qquad \chi^{H^+W^-}_a = \frac{g(H^+W^-h_a)}{M_W/v}, \qquad (1)$$
$$\chi_{abc} = \frac{g(h_ah_bh_c)}{g(hhh)_{SM}}\,.$$

[a]e-mail: ginzburg@math.nsc.ru

The quantities χ^P_a are the ratios of the couplings of h_a with the fundamental particles P to the corresponding couplings for the would be SM Higgs boson with $M_h = M_a$. The other relative couplings describe interaction of h_a with charged Higgs boson. Couplings χ^V_a and $\chi^\pm_a$ are real due to Hermiticity of Lagrangian, while other couplings are generally complex.

(We omit the adjective "relative" below.)

1.1 SM-like scenario

Current data allow us to suggest that Nature realizes the *SM-like scenario*[1]: *The observed particle with mass $M \approx 125$ GeV is a Higgs boson, we call it h_1. It interacts with the gauge bosons and t-quarks with coupling strengths that are close to those predicted by the SM within experimental accuracy (see e.g. [4]-[6]).* In particular, for coupling with the gauge bosons

$$\varepsilon_V = \left|1 - (\chi^V_1)^2\right| \ll 1\,. \qquad (2)$$

1.2 Two Higgs doublet Model (2HDM)

The 2HDM presents the simplest extension of the standard Higgs model [9]. It offers a number of phenomenological scenarios with different physical content in different regions of the model parameter space, such as a natural mechanism for spontaneous CP violation, etc. [9]-[11] For example, the Higgs sector of the MSSM is a particular case of 2HDM. Some variants of 2HDM have interesting cosmological consequences [12],[13].

[1] The term *SM-like scenario* was introduced in [7], the term *alignment limit* was introduced recently for this very situation, see e.g. [8], the *decoupling limit* is the particular case of this scenario.

In the most general 2HDM the couplings (1) obey the following sum rules (SR) [14]-[16]:

$$(a)\ \sum_a (\chi_a^V)^2 = 1, \quad (b)\ |\chi_a^V|^2 + |\chi_a^{H^\pm W^\mp}|^2 = 1, \quad (c)\ \sum_a (\chi_a^f)^2 = 1. \tag{3}$$

We have constructed in [16] the minimal complete set of measurable quantities ("observables") which determine all parameters of the 2HDM. This set contains v.e.v. of Higgs field $v = 246$ GeV, masses of all Higgs bosons $M_{1,2,3}$, $M_\pm$, two out of three couplings χ_a^V, three triple couplings $H^+H^-h_a$ (quantities $\chi_a^\pm$) and one quartic coupling $g(H^+H^-H^+H^-)$. In the most general 2HDM, all these observables are independent of each other. In some particular variants of 2HDM, additional relations between these parameters may appear (for example, in the CP conserving case we have $\chi_3^V = 0$, $\chi_3^\pm = 0$).

1.3 Limitations for parameters

The values of parameters λ_a of 2HDM (and – therefore – mentioned basic parameters) obey two groups of constraints (see e.g. [10], [11]).

The first group is formed by positivity constraints. That are conditions for the stability of Higgs potential at large quasi-classical values of fields. They do not restrict parameters from above.

The second group is formed by perturbativity (and unitarity) constraints. They make it possible to use the first non-vanishing approximation of perturbation theory for description of physical phenomena with reasonable accuracy – *perturbative description*. (This is a tree approximation for most of phenomena and a one-loop approximation for the phenomena which are absent at tree level, e.g. decays $h \to \gamma\gamma$, $h \to Z\gamma$, $h \to gg$). The starting point in obtaining of these constraints is the observation that the effective parameter of perturbative expansion is not λ_i ($i = 1, 2, ...7$) but λ_i/Δ with $\Delta = 8\pi$ or 4π. The perturbativity condition is written usually in the form

$$|\lambda_i| < \Delta\,.$$

At $|\lambda_i| \approx \Delta$ perturbative description of physical phenomena is incorrect even at low energies. In particular, the equations, expressing masses and couplings via parameters of Lagrangian, become invalid. Good example provide the one-loop radiative corrections (RC) to the triple Higgs coupling[2] [17]-[21]. In the SM-like scenario these RC reach $150 \div 200\%$ at $|\lambda_i| \approx \Delta$.

[2]The ref. [21] presents example with clear details. The subject of this paper is Inert Doublet Model, i.e. 2HDM with exact Z_2 symmetry in the SM-like case, at $\lambda_4 = \lambda_5 = 0$ and $\lambda_1 = \lambda_{SM}$. The one-loop corrections to the $g(h_1h_1h_1)$ are described by single parameter λ_3, they reach 180% at $|\lambda_3| \approx \Delta$.

The first non-vanishing approximation of perturbation theory describes physical phenomena with relative inaccuracy k only at

$$|\lambda_i| < k\Delta \qquad (k < 1)\,. \tag{4}$$

In particular, in the region of parameters, provided accuracy of standard description in 30% one should have $k = 0.3$. In this region of parameters the value of RC, discussed in [17]-[21], does not exceed 20%.

Below we will have in mind this very limitation with $k \approx 0.3$.

2 Triple Higgs vertex

The observation of hh production and the extraction of the triple Higgs vertex $g(hhh)$ from the future experiments is scheduled at the LHC and other colliders. This is a necessary step in the verification of the Higgs mechanism. Hopefully, these observations will allow us to see the effects of BSM[3].

The studies of triple Higgs coupling have long history, for recent reviews see e.g. [22]. There are two major parts. The first one is whether it is possible or not to observe hh production, caused by hhh vertex. The second one is whether it is possible to use these observations for extraction of New Physics effect beyond SM.

The accuracy in the extraction of a triple Higgs vertex $g(hhh)$ from the future data cannot be high, since in each case corresponding experiments deal with interference of two channels – an independent production of two Higgses and production of Higgses via hhh vertex. This interference is mainly destructive [23]. For example, for 100 TeV hadron collider with total luminosity $3/abn$ one can hope to reach accuracy of 40% in the extraction of this vertex from future data [24]; at ILC the accuracy in the extraction of $g(hhh)$ will be better than 80% only after 10 years of operation [25]. Therefore, the effects of New Physics will be distinguishable in the data of $g(hhh)$ in the realistic future only if the deviation of this coupling from its SM value is high enough,

$$|\chi_{111} - 1| \gtrsim 1. \tag{5}$$

One of the approaches in the description of the SM violations is to add in the SM Lagrangian terms with anomalous interactions of Higgs boson. It was found for many reasonable benchmark points that these anomalous interactions are difficult for observation [26].

The other approach is to consider some special form of BSM. The review of the whole variety of possible BSM

[3]In the models with additional heavy Higgs bosons h_a with $M_a > 2M_1$ the **resonant** h_1h_1 production like $pp \to (h_2 \to h_1h_1) + ...$ becomes possible. In this paper we discuss only non-resonant h_1h_1 production, without intermediate $h_a \neq h_1$.

models is beyond our scope. We limit ourself considering 2HDM in its most general form.

The potential of such $g(h_1h_1h_1)$ observation was studied for some benchmark points of parameters of 2HDM mainly in the case of CP conservation and with moderate values of parameters [27], for the MSSM with CP-conservation [28] or with violated CP [29] mainly beyond SM-like scenario. The case of SM-like scenario with similar limitations was considered in [30].

2.1 Triple Higgs coupling via observables

The equation for triple Higgs coupling in the most general 2HDM was determined in ref. [16]. We transform it in terms of the introduced observables (see (1)) to the form:

$$
\begin{aligned}
& g(h_1h_1h_1) = (M_1^2/v)\,\chi_{111};\\
& \chi_{111} = \chi_1^V\left[1+\left(1-(\chi_1^V)^2\right)R\right], \quad R = R_1+R_2+R_3;\\
& R_1 = 2\frac{M_\pm^2}{M_1^2}\left(\chi_1^V\chi_1^\pm-1\right), \quad R_2 = 1+2\sum_{b=1}^{3}\frac{M_b^2}{M_1^2}(\chi_b^V)^2, \qquad (6)\\
& R_3 = 2\frac{M_\pm^2}{M_1^2}\left[\sum_{b=2}^{3}\chi_b^V\chi_b^\pm + Re\left(\frac{\chi_1^{H^-W^+}}{\chi_1^V}\sum_{b=1}^{3}\chi_b^{H^+W^-}\chi_b^\pm\right)\right].
\end{aligned}
$$

2.2 Triple Higgs coupling in SM-like scenario

Because of sum rules (3), in the SM-like scenario the couplings of other neutrals h_a with gauge bosons χ_a^V are small. Besides, the absolute value of non-diagonal coupling with EW gauge bosons for the observed Higgs boson $\chi_1^{W^\mp H^\mp}$ is small[4], while similar couplings for the other neutrals $\chi_{2,3}^{W^\pm H^\mp}$ are close to their maximal possible values:

$$\boxed{SR\,(a)} \;\Rightarrow\; |\chi_a^V|^2 < \varepsilon_V \ll 1, \quad a = 2,3. \qquad (7)$$

$$\boxed{SR\,(b)} \;\Rightarrow\; |\chi_1^{H^\pm W^\mp}|^2 \sim \varepsilon_V \ll 1; \quad |\chi_{2,3}^{H^\pm W^\mp}|^2 \approx 1. \qquad (8)$$

With these estimates we have

$$\chi_{111} \approx [1+(R-1/2)\varepsilon_V]. \qquad (9)$$

We see that at moderate values of parameters, the relative coupling χ_{111} is close to 1, and it is difficult to expect a sizable effect[5].

Nevertheless, it is interesting to consider the *special exotic values of model parameters* that provide sizable deviations of triple Higgs coupling from its SM value, i.e. $|\chi_{111}-1| \gtrsim 1$. The perturbativity limitations (4) can suppress some of the discussed opportunities. We consider the effect of different terms R_i, entering R, for the representative case $\varepsilon_V \approx 0.1$.

The term $R_1 = 2(M_\pm/M_1)^2(\chi_1^V\chi_1^\pm - 1)$ can give $|\chi_{111}-1| \gtrsim 1$ if the charged Higgs boson $H^\pm$ is heavy enough and the coupling $\chi_1^\pm$ of the observed Higgs boson with $H^\pm$ deviates substantially from the SM-like value $\chi_1^\pm \approx 1$. If the second condition holds, the two photon width $\Gamma(h_1 \to \gamma\gamma)$ can differ strongly from its SM value. Simultaneous big deviations of triple Higgs coupling and of two photon width from their SM values can be an important source of knowledge about the charged Higgs boson prior to its direct discovery.

The term $R_2 \sim 3+\varepsilon_V(M_b^2/M_1^2)$ can give $|\chi_{111}-1| \gtrsim 1$ only if at least one of other Higgs bosons $h_{2,3}$ is heavy enough, $M_{2,3} > 1$ TeV. Direct discovery of such Higgs bosons seems to be a difficult task. Therefore the value of $g(h_1h_1h_1)$ might become an important source of knowledge about such heavy neutrals for a long time.

The term $R_3 \sim 2(M_\pm/M_1)^2\varepsilon_V\chi_b^\pm$ can give $|\chi_{111}-1| \gtrsim 1$ if the charged Higgs boson is heavy enough and its coupling with one or two heavy neutral Higgs bosons is very large, $\chi_{2,3}^\pm \gtrsim 10$. In this case the two-photon width of such a neutral Higgs boson $h_{2,3}$ will be strongly different from a similar width calculated for the would-be SM Higgs boson with the same mass. At $\chi_{2,3}^\pm \gtrsim 10$, the interaction of charged Higgs bosons can become so strong that a perturbative approach becomes invalid. This case demands special study.

Certainly, the real range of possible values of discussed parameters is restricted by other observations. Better estimates are possible only after measuring of ε_V with reasonable accuracy. In particular, at $\varepsilon_V \ll 0.1$ we cannot expect sizable effects in the triple Higgs vertex.

3 Summary

- Measuring hh production at various colliders is a necessary step in the verification of Higgs mechanism of EWSB. Within the SM-like scenario in the 2HDM, these measurements can give information about the New Physics beyond SM only at exotic values of the parameters listed above. The enlargement of the field of parameters of 2HDM at the transition from CP conserved softly Z_2 broken potential to the most general case gives no new essential opportunities in the deviation of triple Higgs coupling from its SM value.

In our conclusions we limit ourself perturbative limitations in the form (4) with $k \sim 0.3$. These limitations guarantee us applicability of first orders of perturbation theory for description of model (including the expressions of masses and couplings via parameters of Lagrangian) and small value of quantum (loop) corrections.

- In other models deviation of the triple Higgs coupling from its SM value can be stronger than that in 2HDM at

[4]The calculations of $H^- \to W^-h_1$ decay at LHC in [31],[32] are made in the CP-conserving 2HDM and with not very small ε_V.

[5]For the particular CP conserving case and with moderate values of parameters such conclusion was obtained in [30], [27] (see also [28], [29] for the CP conserving MSSM).

moderate values of parameters, see [33]. In the particular case of the nMSSM (2HDM +Higgs singlet) values χ_{111} can range from -5 to 20 [28], [29].

- If the mass M_2 of the heavier Higgs boson h_2 lies within the interval $(250 \div 400)$ GeV and $|\chi_2^t| > 1$ (in the SM-like scenario for h_1), the following interesting phenomenon takes place. The boson h_2 becomes relatively narrow and the cross section of gluon fusion $gg \to h_2$ can be larger than that for the would-be SM Higgs boson with mass M_2. The process $gg \to h_2 \to h_1 h_1$ can be seen as a resonant production of the $h_1 h_1$ pair. In principle, it allows us to discover the mentioned h_2 at LHC (see examples in [27], [34], [35] for special sets of parameters).

Acknowledgements The discussions with F. Boudjema, I.Ivanov, K. Kanishev, M. Krawczyk, P. Osland, M. Vysotsky were useful.

This work was supported in part by grants RFBR 15-02-05868, NCN OPUS 2012/05/B/ST2/03306 (2012-2016) and HARMONIA project under contract UMO-2015/18/M/ST2/00518 (2016-2019).

References

1. ATLAS Collaboration, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett.* **B 716**, 1 (2012); arXiv:1207.7214 [hep-ex]; ATLAS-CONF-2014-010 (2014).
2. CMS Collaboration,*Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC* ; *Phys. Lett.* **B 716**, 30 (2012) arXiv:1207.7235 [hep-ex]; *CMS-PAS- HIG-14-009* (2014).
3. The ATLAS and CMS Collaborations, *Measurements of the Higgs boson production and decay rates and constraints on its couplings from a combined ATLAS and CMS analysis of the LHC pp collision data at* $\sqrt{s} = 7$ *and 8 TeV*, arXiv:1606.02266 [hep-ex]
4. G. Belanger, B. Dumont, U. Ellwanger, J.F. Gunion, S. Kraml, *Higgs Couplings at the End of 2012*, *JHEP* **1302** (2013) 053; arXiv:1212.5244 [hep-ph]; A. Pilaftsis, *Symmetries for SM Alignment in multi-Higgs Doublet Models* arXiv:1602.02017 [hep-ph]
5. B. Dumont, *Higgs Couplings after Moriond*, arXiv:1305.4635 [hep-ph].
6. P.M. Ferreira, R. Guedes, J.F. Gunion, H.E. Haber, M.O.P. Sampaio, R. Santos, *The Wrong Sign limit in the 2HDM*, arXiv:1410.1926 [hep-ph].
7. I.F. Ginzburg, M. Krawczyk, P. Osland,*Distinguishing Higgs Models at Photon Colliders*, 2nd ECFA/DESY Study 1998-2001, 997-1001; hep-ph/9909455;//*Potential of Photon Colliders in resolving SM–like scenarios*, *Nucl. Instrum. Meth.* **A 472** (2001) 149-154; hep-ph/0101229;
   AIP Conf. Proc. Series 578 (2001) 304, IFT-2001-2, 304; *Standard-Model-like scenarios in the 2HDM and Photon Collider potential*, hep-ph/0101331;*Two-Higgs-Doublet Models with CP violation*, Proc. SUSY02 Conference 703-706; hep-ph/0211371.
8. M. Carena, I. Low, N.R. Shah and C.E.M.Wagner, *Impersonating the Standard Model Higgs Boson: Alignment without Decoupling* , *JHEP* 1404 (2014) 015 [arXiv:1310.2248 [hep-ph]]; P.S.B. Dev and A. Pilaftsis, *Maximally Symmetric Two Higgs Doublet Model with Natural Standard Model Alignment*, *JHEP* 1412 (2014) 024; *JHEP* 1511 (2015) 147] [arXiv:1408.3405 [hep-ph]].
9. T.D. Lee, *A Theory of Spontaneous T Violation*, *Phys. Rev.* **D8** (1973) 1226.
10. J.F. Gunion, H.E. Haber, G. Kane, S. Dawson, *The Higgs Hunter's Guide*, (Addison-Wesley, Reading, (1990).
11. G. C. Branco, P. M. Ferreira, L. Lavoura, M. N. Rebelo, M. Sher, J. P. Silva, *Theory and phenomenology of two-Higgs-doublet models*, *Phys. Rept.* **516**, 1 (2012); arXiv:1106.0034 [hep-ph].
12. I. F. Ginzburg, I. P. Ivanov and K. A. Kanishev, *The evolution of vacuum states and phase transitions in 2HDM during cooling of Universe*, *Phys. Rev.* **D 81** (2010) 085031; arXiv:0911.2383 [hep-ph].
13. G. C. Dorsch, S. J. Huber and J. M. No, *A strong electroweak phase transition in the 2HDM after LHC8*, *JHEP* **1310**, 029 (2013); arXiv:1305.6610 [hep-ph].
14. J.F. Gunion, H.E. Haber, J. Wudka, *Sum Rules for Higgs bosons*, *Phys. Rev.* **D 43**, 904 (1991).
15. I.F. Ginzburg, M. Krawczyk, *Symmetries of two Higgs doublet model and CP violation*, *Phys. Rev.* **D 72** (2005) 115013; hep-ph/0408011
16. I.F. Ginzburg, K.A. Kanishev, *2HDM in terms of observable quantities*, *Phys. Rev.* **D 92** (2015) 015024; arXiv:1502.06346 [hep-ph].
17. S. Kanemura, S. Kiyoura, Y. Okada, E. Senaha, C.-P. Yuan, *New Physics Search via the Higgs Self-Coupling*, arXiv:hep-ph/0209326; S. Kanemura, Y. Okada, E. Senaha, C.-P. Yuan, *Higgs coupling constants as a probe of new physics*, arXiv:hep-ph/0211308
18. E. Asakawa, D. Harada, S. Kanemura, Y. Okada, K. Tsumura, *Higgs boson pair production in new physics models at hadron, lepton, and photon colliders*, arXiv:0809.0094; 0902.2458v2 [hep-ph], *Phys.Rev.* **D82** (2010) 115002, arXiv:1009.4670 [hep-ph]
19. P. Osland, P. N. Pandita, L. Selbuz. *Trilinear Higgs couplings in the two Higgs doublet model with CP violation*, arXiv:0802.0060 [hep-ph]
20. F.Boudjema, J.Fujimoto, T.Ishikawa, T.Kaneko, K.Kato, Y.Kurihara, Y.Shimizu, Y.Yasui, *Electroweak Corrections for the Study of the Higgs Potential at the LC*, arXiv:0510184 [hep-ph], F. Boudjema and A. Semenov,*Measurements of the SUSY Higgs self-couplings and the reconstruction of the Higgs potential*, arXiv:hep-ph/0201219
21. A. Arhrib, R. Benbrik, J.El Falaki1z, A. Jueid, *Radiative corrections to the Triple Higgs Coupling in the Inert Higgs Doublet Model*, arXiv:1507.03630 [hep-ph]
22. Y. Yasui, S. Kanemura, S. Kiyoura, K. Odagiri, Okada, E. Senaha, S. Yamashita,*Measurement of the Higgs self-coupling at JLC*, arXiv:hep-ph/0211047 [hep-ph]; G. Panico. *Prospects for double Higgs production*, arXiv:1512.04316 [hep-ph]; J. Baglio, A. Djouadi, R. Grober, M.M. Muhlleitner, J. Quevillon, M. Spira. *The measurement of the Higgs selfcoupling at the LHC: theoretical status*, arXiv:1212.5581 [hep-ph]; J. Baglio. *A theoretical review of triple Higgs coupling studies at the LHC in the Standard Model*, arXiv:1407.1045 [hep-ph]; arXiv:1408.6066 [hep-ph]
23. D.A. Dicus, C. Kao, W.W. Repko, *Interference effects and the use of Higgs boson pair production to study the Higgs trilinear self coupling*, arXiv:1504.02334 [hep-ph]; I.F. Ginzburg, *Higgs boson production at* $\gamma\gamma$ *- colliders (Standard model and beyond)*, Preprint 28(182) Inst. of Mathem., Novosibirsk (1990).
24. A.J. Barr, M.J. Dolan, C. Englert, D.E. Ferreira de Lima, M. Spannowsky, *Higgs Self-Coupling Measurements at a 100 TeV Hadron Collide* , arXiv:1412.7154 [hep-ph]; see also W. Yao *Studies of measuring Higgs self-coupling with* $HH \to bb\gamma\gamma$ *at the future hadron colliders*, arXiv:1308.6302 [hep-ph]; U. Baur. *Measuring the Higgs Boson Self-coupling at High Energy* e^+e^- *Colliders*, arXiv:0906.0028 [hep-ph]

25. T. Barklow, J. Brau, K. Fujii, J. Gao, J. List, N. Walker, K. Yokoya. *ILC Operating Scenarios*, arXiv:1506.07830 [hep-ex]
26. A. Gutierrez-Rodriguez, M. A. Hernandez-Ruz, O.A. Sampayo, A. Chubykalo and A. Espinoza-Garrido. *The Triple Higgs Boson Self-Coupling at Future Linear e^+e^- – Colliders Energies: ILC and CLIC.* arXiv:0807.0663 [hep-ph]; arXiv:0903.1383 [hep-ph]; K. Nishiwaki, S. Niyogi, A. Shivaji, *ttH Anomalous Coupling in Double Higgs Production*, arXiv:1309.6907v2 [hep-ph] F. Goertz, A. Papaefstathiou, L. L. Yang, J. Zurita, *Higgs boson pair production in the D = 6 extension of the SM*, arXiv:1410.3471 [hep-ph]; C.-T. Lu, J. Chang, K. Cheung, J. S. Lee; *An Exploratory study of Higgs-boson pair production.* arXiv:1505.00957 [hep-ph]; H.-J. He, J. Ren, W. Yao, *Probing New Physics of Cubic Higgs Interaction via Higgs Pair Production at Hadron Colliders.* arXiv:1506.03302 [hep-ph]
27. J. Baglio, O. Eberhardt, U. Nierste, M. Wiebusch, *Benchmarks for Higgs Pair Production and Heavy Higgs Searches in the Two-Higgs-Doublet Model of Type II*, *Phys. Rev.* **D 90**, 015008 (2014) arXiv:1403.1264 [hep-ph]; G. Ferrera, J. Guasch, D. Lopez-Val, J. Sola. *Triple Higgs boson production in the Linear Collider.* arXiv:0707.3162 [hep-ph]; A. Arhrib, R. Benbrik, C.-W. Chiang. *Probing triple Higgs couplings of the Two Higgs Doublet Model at Linear Collider.* arXiv:0802.0319 [hep-ph]; A. Arhrib, R. Benbrik, C.-H. Chen, R. Guedes, R. Santos. *Neutral Higgs production in the Two-Higgs doublet model at the LHC.* arXiv:0906.0387 [hep-ph]; F. Cornet, W. Hollik. *Pair Production of Two-Higgs-Doublet Model Light Higgs Bosons in $\gamma\gamma$ Collisions.* arXiv:0808.0719 [hep-ph]; M. Moretti, S. Moretti, F. Piccinini, R. Pittau, J. Rathsman. *Production of Light Higgs Pairs in 2-Higgs Doublet Models via the Higgs-strahlung Process at the LHC.* arXiv:1008.0820v [hep-ph]
28. L. Wu, J.M. Yang, C.-P. Yuan, M. Zhang, *Higgs self-coupling in the MSSM and NMSSM after the LHC Run 1*, arXiv:1504.06932; 1504.07945[hep-ph]; L. Bian, N. Chen, *The Higgs Pair Productions in The CP-violating Two-Higgs-Doublet Model*, arXiv:1607.02703 [hep-ph]
29. J. Cao, D. Li, L. Shang, P. Wu and Y. Zhang, arXiv:1409.8431 [hep-ph]; P. Bandyopadhyay, K. Huitu *Production of two Higgses at the Large Hadron Collider in CP-violating MSSM*, arXiv:1106.5108 [hep-ph]
30. A. Efrati, Y.F. Nir, *What if $g_{hhh} \neq 3m_h^2/v$*, arXiv:1401.0935 [hep-ph].
31. R. Enberg, W. Klemm, S. Moretti, S. Munir, G. Wouda, *Charged Higgs boson in the $W^\pm$ Higgs channel at the Large Hadron Collider* , *Nucl. Phys.* **B893** (2015) 420-442, arXiv:1412.5814[hep-ph].
32. R. Enberg, W. Klemm, S. Moretti, S. Munir, G. Wouda, *Search for Charged Higgs bosons via decays to $W^\pm$ and a 125 GeV Higgs at the Large Hadron Collider*, arXiv:1502.02931 [hep-ph]
33. S.I. Godunov, A.N. Rozanov, M.I. Vysotsky, E.V. Zhemchugov,*Extending the Higgs sector: an extra singlet* , arXiv:1503.01618 (2015); *Double Higgs boson production in the Standard Model with extra scalar particles*, E.V. Zhemchugov, arXiv:1510.02382 [hep-ph]
34. V. Barger, L.L. Everett, C.B. Jackson, A.D. Peterson, G. Shaughnessy,*Measuring the 2HDM Scalar Potential at LHC14*, *Phys. Rev.* **D 90**, 095006 (2014) arXiv:1408.2525 [hep-ph].
35. S. Kanemura, H. Yokoya, Y.-J. Zheng, *Searches for additional Higgs bosons in multi-top-quarks events at the LHC and the International Linear Collider*, arXiv:1505.01089 [hep-ph].